

Giuseppe Manganelli*

TWO ADDITIONAL RECORDS FOR *LEIOSTYLA* IN TURKEY
(PULMONATA, ORCULIDAE/PUPILLIDAE)**

KEY WORDS: *Leiostyla*, Turkey.

Abstract

Two species of the genus *Leiostyla*, *L. tenuimarginata* (PILSBRY) and *L. sinangula* (SCHILEYKO), are reported for the first time from Turkey.

Riassunto:

Tra i materiali raccolti negli anni trascorsi in Turchia da alcuni zoologi italiani sono state rinvenute alcune conchiglie riferibili a due specie del genere *Leiostyla*: *L. tenuimarginata* (PILSBRY) e *L. sinangula* (SCHILEYKO), mai segnalate prima in Turchia anche se conosciute per la vicina Georgia (USSR).

Many Italian zoologists have visited Turkey in the last twenty years and collected molluscs which were given to Prof. GIUSTI of the Dept. of Evolutionary Biology of the University of Siena.

Among these materials two lots of *Leiostyla* were found. *Leiostyla* is a genus of the Lauriinae (Pulmonata: Orculidae according to PILSBRY, 1922; SCHILEYKO 1984; or Pupillidae ecc. to ZILCH, 1959, 1985; SCHILEYKO, 1975; BACKHUYSEN, 1975; DAMJANOV & LIKHAREV, 1975; KERNEY et al., 1983; WALDEN, 1983; HOLYOAK & SEDDON, 1986; GITTEMBERGER & PIEPER, 1988; DE WINTER & RIPKEN, 1989) with a broad discontinuous distribution in the Palearctic. It inhabits the Macaronesian Is., Maghreb and Atlantic Europe (Ireland, Great Britain, France, Spain, Portugal), is absent from central southern Europe and reappears in Bulgaria, Turkey, Crimea, Caucasus and North Iran. In the oriental sector, 11-13 species have been reported most from the Caucasus Region; in other areas only presences of apparently relict type have been documented (cf. MANGANELLI et al., 1990).

The first record of *Leiostyla* in Turkey was published by RETOWSKY (1889) who described *Pupa (Charadrobia) pontica* from Samsum. This species was regarded as valid by PILSBRY (1922) and ZILCH (1985) and as a junior synonym of *Leiostyla pulchra* (RETOWSKY, 1883) by LIKHAREV & RAMMEL'MEIER (1952) and SCHILEYKO (1975, 1984). Later LINDHOLM (1914)

(*) Dipartimento di Biologia Evolutiva, Università di Siena, Via Mattioli 4, I-53100 Siena, Italy.

(**) Research financed by a CNR «Gruppo di Biologia Naturalistica» and MPI 40% and 60% grants.

(***) Lavoro accettato il 6 aprile 1990.

described *Lauria paulinae* subsp. *unicolumellaris* from Mt. Salolet at Artvin. This taxon was regarded by SCHILEYKO (1975, 1984) as a simple form of *L. paulinae* (LINDHOLM, 1913). Many years later a third record was published by GOTTING (1963) of a new species named *Leiostyla schweigeri* from Abant-Gebirge near Bolu.

Description of two species from Turkey and a third from the Russian - Turkish border are in press (B. POKRYSZKO pers. com.).

The two lots of materials in my possession correspond to two Russian (Georgian) species never before reported in Turkey:

Leiostyla tenuimarginata (PILSBRY, 1922)

Lauria (*Leiostyla*) *tenuimarginata* PILSBRY, 1922: 76-77, 1923: Pl. 10, figs. 1-2.

Euxinolauria (*Caucasipupa*) *tenuimarginata*, SCHILEYKO, 1984: 137-138, Figs. 64-65.

Material examined

Turkey: Borçka (Artvin), G. OSELLA leg. 18.VI.1969 (1 sp.) GIUSTI collection (University of Siena, Italy).

Leiostyla sinangula (SCHILEYKO, 1975)

Euxinolauria (*Caucasipupa*) *sinangula* SCHILEYKO, 1975: 1775, Figs. 3,3-4, 6,7.

Euxinolauria (*Caucasipupa*) *sinangula*, SCHILEYKO, 1984: 143-145, Figs. 72-73.

Material examined

Turkey: Borçka (Artvin), V. COTTARELLI leg. 15-18.VI.1969 (2 sps.) GIUSTI collection (University of Siena, Italy).

Comments

The present report adds two more *Leiostyla* species to the malacofauna of Turkey: *L. tenuimarginata* and *L. sinangula*. This however is not very surprising since Borçka (Artvin) is close to the three Georgian (USSR) sites where the two species are known.

L. tenuimarginata was known from Notanebi Post, 40 km north of Batumi (type locality); Botanical Gardens of Batumi; Ciakvistavi, 20 km east of Batumi (LIKHAREV & RAMMEL'MEIER, 1952; SCHILEYKO, 1975, 1984). Although our specimen (Fig. 1) is more cylindrical (less conical) in shape than those reported in the literature (cf. PILSBRY, 1923: Pl. 10, figs. 1-2; LIKHAREV




Fig. 1. *Leiostyla tenuimarginata* (PILSBRY, 1922). Borçka (Artvin, Turkey), G. OSELLA leg. 18.VI.1969.

Fig. 2. *Leiostyla sinangula* (SCHILEYKO, 1984). Borçka (Artvin, Turkey), V. COTTARELLI leg. 15-18.VI.1969.

& RAMMEL'MEIER, 1952; Fig. 67; SCHILEYKO, 1984: 68) it corresponds very well to the latter and to a shell examined by us (SMF 52113) in the simplified structure of the apertural armature. This character makes *L. tenuimarginata* one of the most peculiar oriental *Leiostyla* species.

The other species was also reported from the Botanical Gardens of Batumi where it was found together with *L. tenuimarginata* in the litter and on bamboo stems. The only adult specimen in my possession resembles that illustrated by SCHILEYKO (1984: Fig. 72) in shape, in the presence of two horizontal columellar lamellae, and in the form of the peristomial knob and crenulated peristome. These characters also correspond to those of the shell figured and erroneously (SCHILEYKO, *pers. com.*) referred to as *L. paulinae* by LIKHAREV & RAMMEL'MEIER (1952: Fig. 66).

Acknowledgments

I wish to thank Prof. F. GIUSTI, Prof. A.A. SCHILEYKO and Dr. B. POKRYSZKO for the information and advice. I also wish to thank Mr. L. GAMBERUCCI and Mrs. A. DAVIDDI for technical assistance.

R E F E R E N C E S

- BACKHUYSEN, W. (1975). Zoogeography and taxonomy of the land and freshwater molluscs of the Azores. Backhuys & Meesters, Amsterdam, 350 pags., 97 maps, 32 Pls.
- DAMJANOV, S.G. & LIKHAREV, I. (1975). Gastropoda terrestria. *Fauna bulgarica*, **4**: 425 pags.
- DE WINTER, A.J. & RIPKEN, Th. E.J. (1988). *Leiostyla anglica* (WOOD) on Madeira (Gastropoda Pulmonata, Pupillidae). *Basteria*, **53**: 47-48.
- GITTENBERGER, E. & PIEPER, H. (1988). Eine neue *Leiostyla* - Arten aus dem Iran, (Pulmonata: Pupillidae). *Arch. Molluskenk.*, **119**: 65-71.
- GÖTTING, K.J. (1963). *Leiostyla schweigeri* n. sp. und *Pseudamnicola lindbergi* BOETTGER - zwei interessante Gastropoden aus Anatolien. *Arch. Molluskenk.*, **92**: 31-33.
- HOLYOAK, D.T. & SEDDON, M.B. (1986). An undescribed *Leiostyla* (Gastropoda: Pupillidae) from Madeira. *J. Conchol.*, **32**: 191-193.
- KERNEY, M.P., CAMERON R.A.D. & JUNGBLUTH, J.H. (1983). Die Landschnecken Nord - und Mitteleuropas. Paul Parey, Hamburg und Berlin, 384 pags., 24 Pls.
- LIKHAREV, I.M. & RAMMEL'MEIER, E.S. (1952). Terrestrial mollusks of the Fauna of the U.S.S.R. English translation, Monson, Jerusalem, 574 pagg.
- LINDHOLM, W.A. (1913). Beschreibung neuer Arten und Formen aus dem Kaukasus-Gebiete. *Nachrichtenbl. dtsc. malakozool. Ges.*, **45**: 17-23, 62-69.
- LINDHOLM, W.A. (1914). Beschreibung vier neuer Landschnecken und einer neuen Untergattung aus dem sudwestlichen Transkaukasien. *Nachrichtenbl. dtsc. malakozool. Ges.*, **46**: 33-38.
- MANGANELLI, G.; GIUSTI, F & DELLE CAVE, L. (1990). Notulae Malacologicae, XLVIII. Lauriinae (Gastropoda: Pulmonata, Oculidae/Pupillidae) from the Villafranchian of peninsular Italy. *Basteria*, **53**, in press.
- PILSBRY, H.A., 1922-1923. Manual of Conchology - Structural and systematic. With illustrations of the species. (II) 27 (105): 1-80, Pls. 1-5 (1922); 27 (106): 81-128, Pls. 6-11 (1923).
- ROTOWSKY, O. (1889). Liste der von mir auf meiner Reise von Kostantinopel nach Batum gesammelten Binnenmollusken. *Ber. Senckenb. naturforsch. Ges.*, **1889**: 225-265.
- SCHILEYKO, A.A. (1975). Molluscs of the subfamily Lauriinae (Pulmonata, Pupillidae) of the USSR. *Zool. Zh.*, **54**: 1767-1782. (in Russian).
- SCHILEYKO, A.A. (1984). Molluscs. Terrestrial molluscs of the suborder Pupillina of the fauna of the URSS (Gastropoda, Pulmonata, Geophila). *Fauna SSSR*, (N.S.), **130**: 1-399. (in Russian).
- WALDEN, H.W. (1983). Systematic and biogeographical studies of the terrestrial Gastropoda of Madeira. With an annotated check-list. *Ann. zool. fenn.*, **20**: 255-275.
- ZILCH, A. (1959). Gastropoda. Euthyneura. *Handb. Paläzoöl* 6 (2, 2nd part): I-XII + 1-200.
- ZILCH, A. (1985). Die Typen und Typoide des Natur-Museums Senckenberg, 75: Mollusca, Pupillacea (5): Pupillidae: Lauriinae, Argininae. *Arch. Molluskenk.*, **116**: 119-136.